

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/64, 31/44, 31/155 // (A61K 31/64, 31:44, 31:155)	A1	(11) International Publication Number: WO 99/03477 (43) International Publication Date: 28 January 1999 (28.01.99)
(21) International Application Number: PCT/GB98/02110 (22) International Filing Date: 16 July 1998 (16.07.98) (30) Priority Data: 9715295.3 18 July 1997 (18.07.97) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): BUCKINGHAM, Robin, Edwin [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). SMITH, Stephen, Alistair [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Beecham Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). (74) Agent: RUTTER, Keith; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: TREATMENT OF DIABETES WITH THIAZOLIDINEDIONE, INSULIN SECRETAGOGUE AND DIGUANIDE (57) Abstract A method for the treatment of diabetes mellitus and conditions associated with diabetes mellitus in a mammal, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, an insulin secretagogue and a biguanide antihyperglycaemic agent, to a mammal in need thereof; and composition for use in such method.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TREATMENT OF DIABETES WITH THIAZOLIDINEDIONE, INSULIN SECRETAGOGUE AND DIGUANIDE

This invention relates to a method of treatment, in particular to a method for the treatment of diabetes mellitus, especially non-insulin dependent diabetes
5 (NIDDM) (or Type 2 diabetes) and conditions associated with diabetes mellitus.

Insulin secretagogues are compounds which promote increased secretion of insulin by the pancreatic beta cells.

The sulphonylureas are well known examples of insulin secretagogues. The sulphonylureas act as antihyperglycaemic agents and are used in the treatment of Type
10 2 diabetes). Examples of sulphonylureas include glibenclamide, glipizide, gliclazide, glimepiride, tolazamide and tolbutamide.

Biguanide antihyperglycaemic agents are commonly used in the treatment of Type 2 diabetes). 1,1 - Dimethylbiguanidine (or Metformin) is an example of a biguanide antihyperglycaemic agent.

15 European Patent Application, Publication Number 0,306,228 relates to certain thiazolidinedione derivatives disclosed as having antihyperglycaemic and antihyperlipidaemic activity. One particular thiazolidinedione disclosed in EP 0306228 is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (hereinafter 'Compound (I)'). WO94/05659 discloses certain salts of Compound
20 (I) including the maleate salt.

Compound (I) is an example of a class of anti-hyperglycaemic agents known as 'insulin sensitisers'. In particular Compound (I) is a thiazolidinedione insulin sensitiser.

European Patent Applications, Publication Numbers: 0008203, 0139421,
25 0032128, 0428312, 0489663, 0155845, 0257781, 0208420, 0177353, 0319189, 0332331, 0332332, 0528734, 0508740; International Patent Application, Publication Numbers 92/18501, 93/02079, 93/22445 and United States Patent Numbers 5104888 and 5478852, also disclose certain thiazolidinedione insulin sensitisers.

Another series of compounds generally recognised as having insulin sensitiser
30 activity are those typified by the compounds disclosed in International Patent Applications, Publication Numbers WO93/21166 and WO94/01420. These compounds are herein referred to as 'acyclic insulin sensitisers'. Other examples of acyclic insulin sensitisers are those disclosed in United States Patent Number 5232945 and International Patent Applications, Publication Numbers WO92/03425 and
35 WO91/19702.

Examples of other insulin sensitisers are those disclosed in European Patent Application, Publication Number 0533933, Japanese Patent Application Publication Number 05271204 and United States Patent Number 5264451.

The above mentioned publications are incorporated herein by reference.

It is now surprisingly indicated that Compound (I) in combination with an insulin secretagogue and a biguanide antihyperglycaemic agent provides a particularly beneficial effect on glycaemic control, such combination is therefore particularly
 5 useful for the treatment of diabetes mellitus, especially Type 2 diabetes and conditions associated with diabetes mellitus. The treatment is also indicated to proceed with minimum side effects.

Accordingly, the invention provides a method for the treatment of diabetes mellitus, especially Type 2 diabetes, and conditions associated with diabetes mellitus,
 10 in a mammal such as a human, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, an insulin secretagogue and a biguanide antihyperglycaemic agent, to a mammal in need thereof.

The method comprises either co-administration of the insulin sensitiser, an insulin secretagogue and a biguanide antihyperglycaemic agent or sequential
 15 administration thereof.

Co-administration includes administration of a formulation which includes an insulin sensitiser, an insulin secretagogue and a biguanide antihyperglycaemic agent or the essentially simultaneous administration of separate formulations of each agent.

In another aspect the invention provides the use of an insulin sensitiser, such
 20 as Compound (I), an insulin secretagogue and a biguanide antihyperglycaemic agent, in the manufacture of a composition for the treatment of diabetes mellitus, especially Type 2 diabetes and conditions associated with diabetes mellitus.

A suitable insulin sensitiser is a thiazolidinedione insulin sensitiser.

A suitable thiazolidinedione insulin sensitiser is Compound (I).

Other suitable thiazolidinedione insulin sensitisers include (+) -5-[[4-[(3,4-
 25 dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl)methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl] thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl] thiazolidine-2,4-dione (or pioglitazone) or 5-[(2-
 30 benzyl-2,3-dihydrobenzopyran)-5-ylmethyl]thiazolidine-2,4-dione (or englitazone)

Suitable insulin secretagogues include sulphonylureas.

Suitable sulphonylureas include glibenclamide, glipizide, gliclazide, glimepiride, tolazamide and tolbutamide.

Further sulphonylureas include acetohexamide, carbutamide,
 35 chlorpropamide, glibornuride, gliquidone, glisentide, glisolamide, glisoxepide, glycopyramide and glycylamide.

Further suitable insulin secretagogues include repaglinide.

A suitable biguanide antihyperglycaemic agent is metformin, buformin or phenformin, especially metformin.

In one particular aspect, the method comprises the administration of 2 to 12 mg of Compound (I), especially when administered per day.

5 Particularly, the method comprises the administration of 2 to 4, 4 to 8 or 8 to 12 mg of Compound (I) per day.

Particularly, the method comprises the administration of 2 to 4mg of Compound (I), especially when administered per day.

10 Particularly, the method comprises the administration of 4 to 8mg of Compound (I), especially when administered per day.

Particularly, the method comprises the administration of 8 to 12 mg of Compound (I), especially when administered per day.

Preferably, the method comprises the administration of 2 mg of Compound (I), especially when administered per day.

15 Preferably, the method comprises the administration of 4 mg of Compound (I), especially when administered per day.

Preferably, the method comprises the administration of 8 mg of Compound (I), especially when administered per day.

20 It will be understood that the insulin sensitiser, such as compound (I), the insulin secretagogue and the biguanide antihyperglycaemic agent are each administered in a pharmaceutically acceptable form, including pharmaceutically acceptable derivatives such as pharmaceutically acceptable salts, esters and solvates thereof, as appropriate. In certain instances herein the names used for the relevant insulin secretagogues and the biguanide antihyperglycaemic agents may relate to a
25 particular pharmaceutical form of the relevant active agent: It will be understood that all pharmaceutically acceptable forms of the active agents per se are encompassed by this invention.

Suitable pharmaceutically acceptable salted forms of the insulin sensitisers, such as Compound (I), include those described in the above mentioned patents and
30 patent applications such as in EP 0306228 and WO94/05659 for Compound (I). A preferred pharmaceutically acceptable salt for Compound (I) is a maleate.

Suitable pharmaceutically acceptable solvated forms of the insulin sensitisers, such as Compound (I), include those described in the above mentioned patents and patent applications, such as in EP 0306228 and WO94/05659 for
35 Compound (I), in particular hydrates.

Suitable pharmaceutically acceptable forms of the insulin secretagogue and the biguanide antihyperglycaemic agent depend upon the particular compound used but include known pharmaceutically acceptable forms of the particular compound

chosen. Such derivatives are found or are referred to in standard reference texts such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) (for example see the 31st Edition page 341 and pages cited therein).

5 The insulin sensitisers, such as Compound (I) or, a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, may be prepared using known methods, for example those disclosed in the above mentioned patents and patent applications, such as EP 0306228 and WO94/05659 for Compound (I). The disclosures of the above mentioned patents and patent applications, such as
10 EP 0306228 and WO94/05659, are incorporated herein by reference.

 Compound (I) may exist in one of several tautomeric forms, all of which are encompassed by the term Compound (I) as individual tautomeric forms or as mixtures thereof. Compound (I) contains a chiral carbon atom, and hence can exist in up to two stereoisomeric forms, the term Compound (I) encompasses all of these isomeric forms
15 whether as individual isomers or as mixtures of isomers, including racemates.

 The insulin secretagogue and biguanide antihyperglycaemic agent of choice is prepared according to known methods, such methods are found or are referred to in standard reference texts, such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra
20 Pharmacopoeia (London, The Pharmaceutical Press) (for example see the 31st Edition page 341 and pages cited therein).

 When used herein the term 'conditions associated with diabetes' includes conditions associated with diabetes mellitus itself and complications associated with diabetes mellitus. Also included in 'conditions associated with diabetes' are those
25 conditions associated with the pre-diabetic state.

 When used herein the term 'conditions associated with the pre-diabetic state' includes conditions such as insulin resistance, including hereditary insulin resistance, impaired glucose tolerance and hyperinsulinaemia.

 'Conditions associated with diabetes mellitus itself' include hyperglycaemia,
30 insulin resistance, including acquired insulin resistance. Further conditions associated with diabetes mellitus itself include hypertension and cardiovascular disease, especially atherosclerosis and conditions associated with insulin resistance. Conditions associated with insulin resistance include polycystic ovarian syndrome and steroid induced insulin resistance and gestational diabetes.

35 'Complications associated with diabetes mellitus' includes renal disease, especially renal disease associated with Type 2 diabetes, neuropathy and retinopathy.

 Renal diseases associated with Type 2 diabetes include nephropathy, glomerulonephritis, glomerular sclerosis, hypertensive nephrosclerosis and end stage

renal disease. Additional renal diseases associated with Type 2 diabetes include nephrotic syndrome.

For the avoidance of doubt, when reference is made herein to scalar amounts, including mg amounts, of Compound (I) in a pharmaceutically acceptable form, the scalar amount referred to is made in respect of Compound (I) *per se*: For example 2 mg of Compound (I) in the form of the maleate salt is that amount of maleate salt which contains 2 mg of Compound (I).

Diabetes mellitus is preferably Type 2 diabetes.

The particularly beneficial effect on glycaemic control provided by the treatment of the invention is indicated to be a synergistic effect relative to the control expected for the sum of the effects of the individual active agents.

Suitably the insulin sensitiser is the agent of first administration.

Glycaemic control may be characterised using conventional methods, for example by measurement of a typically used index of glycaemic control such as fasting plasma glucose or glycosylated haemoglobin (HbA1c). Such indices are determined using standard methodology, for example those described in: Tiescher A, Richterich, P., Schweiz. med. Wschr. 101 (1971), 345 and 390 and Frank P., 'Monitoring the Diabetic Patient with Glycosolated Hemoglobin Measurements', Clinical Products 1988

In a preferred aspect, the dosage level of each of the active agents when used in accordance with the treatment of the invention will be less than would have been required from a purely additive effect upon glycaemic control.

There is also an indication that the treatment of the invention will effect an improvement, relative to the individual agents, in the levels of advanced glycosylation end products (AGEs), leptin and serum lipids including total cholesterol, HDL-cholesterol, LDL-cholesterol including improvements in the ratios thereof, in particular an improvement in serum lipids including total cholesterol, HDL-cholesterol, LDL-cholesterol including improvements in the ratios thereof.

As used herein the term 'pharmaceutically acceptable' embraces both human and veterinary use: for example the term 'pharmaceutically acceptable' embraces a veterinarily acceptable compound.

In the method of the invention, the active medicaments are preferably administered in pharmaceutical composition form. As indicated above, such compositions can include all medicaments or one only of the medicaments.

Accordingly, in one aspect the present invention also provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and a biguanide antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor.

Such compositions may be prepared by admixing an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and a biguanide antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor.

5 Usually the compositions are adapted for oral administration. However, they may be adapted for other modes of administration, for example parenteral administration, sublingual or transdermal administration.

 The compositions may be in the form of tablets, capsules, powders, granules, lozenges, suppositories, reconstitutable powders, or liquid preparations, such as oral
10 or sterile parenteral solutions or suspensions.

 In order to obtain consistency of administration it is preferred that a composition of the invention is in the form of a unit dose.

 Unit dose presentation forms for oral administration may be tablets and capsules and may contain conventional excipients such as binding agents, for example
15 syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate; disintegrants, for example starch, polyvinylpyrrolidone, sodium starch glycollate or microcrystalline cellulose; or pharmaceutically acceptable wetting agents such as sodium lauryl sulphate.

20 The compositions are preferably in a unit dosage form in an amount appropriate for the relevant daily dosage.

 Suitable dosages for the insulin sensitisers include those disclosed in the abovementioned patents and patent applications.

 Suitable dosages, including unit dosages, of Compound (I) comprise 1, 2, 3,
25 4, 5, 6, 7, 8, 9, 10, 11 or 12 mg of Compound (I).

 In the treatment the medicaments may be administered from 1 to 6 times a day, but most preferably 1 or 2 times per day.

 Particular dosages of Compound (I) are 2mg/day, 4mg/day, including 2mg twice per day, and 8 mg/day, including 4mg twice per day.

30 Suitable dosages including unit dosages of the insulin secretagogue, such as the sulphonylurea, or the biguanide antihyperglycaemic agent, include the known dosages including unit doses for these compounds as described or referred to in reference text such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra
35 Pharmacopoeia (London, The Pharmaceutical Press) (for example see the 31st Edition page 341 and pages cited therein).

 Thus for the sulphonylureas, a typical daily dosage of glibenclamide is in the range of from 2.5 to 20 mg, for example 10mg twice per day or 20mg once per day; a

typical daily dosage of glipizide is in the range of from 2.5 to 40 mg; a typical daily dosage of gliclazide is in the range of from 40 to 320 mg; a typical daily dosage of tolazamide is in the range of from 100 to 1000 mg; a typical daily dosage of tolbutamide is in the range of from 1000 to 3000 mg; a typical daily dosage of
5 chlorpropamide is in the range of from 100 to 500 mg; and a typical daily dosage of gliquidone is in the range of from 15 to 180 mg.

Repaglinide may be taken in amounts, usually in the range of from 0.5mg to 4mg and usually with meals, up to a typical maximum daily dosage of 16mg per day.

With regard to the biguanide antihyperglycaemic agents, suitable dosages of
10 metformin include up to 3000mg per day, in unit doses of 500mg (for example two or three times per day) or 850mg (for example two times per day), one example of a dosage for metformin is 500mg once building to five times per day.

The solid oral compositions may be prepared by conventional methods of blending, filling or tableting. Repeated blending operations may be used to distribute
15 the active agent throughout those compositions employing large quantities of fillers. Such operations are of course conventional in the art. The tablets may be coated according to methods well known in normal pharmaceutical practice, in particular with an enteric coating.

Oral liquid preparations may be in the form of, for example, emulsions,
20 syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel, hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan
25 monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid; and if desired conventional flavouring or colouring agents.

30 For parenteral administration, fluid unit dosage forms are prepared utilizing the compound and a sterile vehicle, and, depending on the concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the compound can be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, a
35 preservative and buffering agents can be dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the Compound (I) is suspended in the vehicle instead of being

dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

5 Compositions may contain from 0.1% to 99% by weight, preferably from 10-60% by weight, of the active material, depending upon the method of administration.

 Composition may, if desired, be in the form of a pack accompanied by written or printed instructions for use.

10 The compositions are prepared and formulated according to conventional methods, such as those disclosed in standard reference texts, for example the British and US Pharmacopoeias, 's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) and Harry's Cosmeticology (Leonard Hill Books) (for example see the 31st Edition page
15 341 and pages cited therein).

 The present invention also provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and a biguanide antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor, for use as an active therapeutic
20 substance.

 The invention also provides the use of an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and a biguanide antihyperglycaemic agent for the manufacture of a medicament for the treatment of diabetes mellitus and conditions associated with diabetes.

25 In particular, the present invention provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and a biguanide antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor, for use in the treatment of diabetes mellitus and conditions associated with diabetes mellitus.

30 A range of 2 to 4mg includes a range of 2.1 to 4, 2.2 to 4, 2.3 to 4, 2.4 to 4, 2.5 to 4, 2.6 to 4, 2.7 to 4, 2.8 to 4, 2.9 to 4 or 3 to 4mg.

 A range of 4 to 8mg includes a range of 4.1 to 8, 4.2 to 8, 4.3 to 8, 4.4 to 8, 4.5 to 8, 4.6 to 8, 4.7 to 8, 4.8 to 8, 4.9 to 8, 5 to 8, 6 to 8 or 7 to 8mg.

 A range of 8 to 12 mg includes a range of 8.1 to 12, 8.2 to 12, 8.3 to 12, 8.4
35 to 12, 8.5 to 12, 8.6 to 12, 8.7 to 12, 8.8 to 12, 8.9 to 12, 9 to 12, 10 to 12 or 11 to 12mg.

 No adverse toxicological effects are expected for the compositions or methods of the invention in the abovementioned dosage ranges.

Composition for compound (I)

Preparation of Concentrate: Tableting concentrate was prepared using the following materials

Ingredient	Quantity (%)
Milled Compound (I) as maleate salt	13.25 (pure maleate salt)
Sodium Starch Glycollate	5.00
Hydroxypropyl Methylcellulose 2910	5.00
Microcrystalline Cellulose (Avicel PH102)	20.0
Lactose Monohydrate, regular grade	to 100
Purified water	*

* Removed during processing.

The concentrate was then formulated into tablets using the following:

Tablet Strength	Quantity (mg per Tablet)			
	1.0mg	2.0mg	4.0mg	8.0mg
Active Ingredient:				
Compound (I) maleate Concentrate granules	10.00	20.00	40.00	80.00
Other Ingredients:				
Sodium Starch Glycollate	6.96	6.46	5.46	10.92
Microcrystalline Cellulose (Avicel PH102)	27.85	25.85	21.85	43.70
Lactose monohydrate, (Pharmatose DCL15),	104.44	96.94	81.94	163.88
Magnesium Stearate	0.75	0.75	0.75	1.50
Total Weight of Tablet Core	150.0	150.0	150.0	300.0
Opadry	4.5	4.5	4.5	9.0
Total Weight of Film Coated Tablet	154.5	154.5	154.5	309.0

Compositions for other active agents are as described in the above mentioned publications.

Claims

1. A method for the treatment of diabetes mellitus and conditions associated with diabetes mellitus in a mammal, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, an insulin secretagogue and a biguanide antihyperglycaemic agent, to a mammal in need thereof.
2. A method according to claim 1, wherein the insulin secretagogue is glibenclamide, glipizide, gliclazide, glimepiride, tolazamide, tolbutamide, acetohexamide, carbutamide, chlorpropamide, glibornuride, gliquidone, glisentide, glisolamide, glisoxepide, glyclopyamide, glycylamide or repaglinide.
3. A method according to claim 1, wherein the biguanide is metformin, buformin or phenformin.
4. A method according to any one of claims 1 to 3, wherein the insulin sensitiser is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (Compound I).
5. A method according to any one of claims 1 to 4, which comprises the administration of 2 to 12 mg of Compound (I).
6. A method according to any one of claims 1 to 5, which comprises the administration of 2 to 4, 4 to 8 or 8 to 12 mg of Compound (I).
7. A method according to any one of claims 1 to 6, which comprises the administration of 2 to 4mg of Compound (I).
8. A method according to any one of claims 1 to 6, which comprises the method the administration of 4 to 8mg of Compound (I).
9. A method according to any one of claims 1 to 6, which comprises the administration of 8 to 12 mg of Compound (I).
10. A method according to any one of claims 1 to 6, which comprises the administration of 2 mg of Compound (I).

11. A method according to any one of claims 1 to 6, which comprises the administration of 4 mg of Compound (I).
12. A method according to any one of claims 1 to 6, which comprises the
5 administration of 8 mg of Compound (I).
13. A method according to claim 1, wherein the insulin sensitiser is (+) -5-[[4-[(3,4-dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl] thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl] thiazolidine-2,4-dione (or pioglitazone) or
10 5-[(2-benzyl-2,3-dihydrobenzopyran)-5-ylmethyl]thiazolidine-2,4-dione (or englitazone); or a pharmaceutically acceptable form thereof.
14. A pharmaceutical composition comprising an insulin sensitiser, an insulin secretagogue, a biguanide antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor.
15. A composition according to claim 14, wherein the insulin secretagogue is a
20 sulphonylurea.
16. A composition according to claim 14 or claim 15, wherein the insulin secretagogue is glibenclamide, glipizide, gliclazide, glimepiride, tolazamide or tolbutamide, acetohexamide, carbutamide, chlorpropamide, glibornuride,
25 gliquidone, glisentide, glisolamide, glisoxepide, glyclopyamide, glycyclamide or repaglinide.
17. A composition according to claim 1, wherein the biguanide is metformin, buformin or phenformin.
30
18. A composition according to any one of claims 14 to 17, wherein the insulin sensitiser is Compound (I)
19. A composition according to any one of claims 14 to 17, which comprises 2 to 12
35 mg of Compound (I).

20. A pharmaceutical composition comprising an insulin sensitiser an insulin secretagogue and a pharmaceutically acceptable carrier therefor, for use as an active therapeutic substance.
- 5 21. A pharmaceutical composition comprising an insulin sensitiser, an insulin secretagogue, a biguanide antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor, for use in the treatment of diabetes mellitus and conditions associated with diabetes mellitus.
- 10 22. A composition according to any one of claims 14, 20 or 21, wherein the insulin sensitiser is (+) -5-[[4-[(3,4-dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl] thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl] thiazolidine-2,4-dione (or pioglitazone) or 5-[(2-benzyl-2,3-dihydrobenzopyran)-5-ylmethyl]thiazolidine-2,4-dione (or englitazone); or a pharmaceutically acceptable form thereof.
- 15

INTERNATIONAL SEARCH REPORT

Internat. Application No

PCT/GB 98/02110

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/64 A61K31/44 A61K31/155 //(A61K31/64,31:44,31:155)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 749 751 A (TAKEDA CHEMICAL INDUSTRIES LTD) 27 December 1996 see page 9, line 42-55; claims 24-26; example 2 see page 8, line 32-34	1-22
Y,P	KAPPEL C. ET AL: "Type 2 diabetes: Update on therapy" COMPREHENSIVE THERAPY, vol. 24, no. 6-7, June 1998, pages 319-326, XP002080974 see page 324, column 2, paragraph 2; figure 1	1-22



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 October 1998

Date of mailing of the international search report

02/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gonzalez Ramon, N

INTERNATIONAL SEARCH REPORT

Intern. Application No.

PCT/GB 98/02110

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ELSON DF ET AL: "Therapy for type 2 diabetes mellitus 'see comments!'" WMJ, MAR 1998, 97 (3) P49-54, XP002081122 UNITED STATES see page 52	1-22
Y	see page 53, column 1, paragraph 2 ---	1-22
A	SCHEEN AJ ET AL: "Oral antidiabetic agents. A guide to selection." DRUGS, FEB 1998, 55 (2) P225-36, XP002080821 NEW ZEALAND see page 232, column 2, paragraph 1; figure 1 ---	1-22
A	WO 93 03724 A (UPJOHN CO) 4 March 1993 see claims 3,4,10,11 ---	1-22
A	WO 96 09823 A (SHAMAN PHARMACEUTICALS INC) 4 April 1996 see claims 7-11 ---	1-22
A	WOLFFENBUTTEL B. ET AL: "New treatments for patients with type 2 diabetes mellitus" POSTGRAD MED, vol. 72, pages 657-662, XP002081123 boxes 3,4,5 see abstract -----	1-22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 98/02110

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 1-13
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/02110

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0749751 A	27-12-1996	AU 5603496 A	09-01-1997
		CA 2179584 A	21-12-1996
		CN 1145783 A	26-03-1997
		CZ 9601811 A	15-01-1997
		EP 0861666 A	02-09-1998
		HU 9601698 A	28-05-1997
		JP 9067271 A	11-03-1997
		JP 10167986 A	23-06-1998
		NO 962606 A	23-12-1996
		SK 79496 A	08-01-1997
WO 9303724 A	04-03-1993	AT 151634 T	15-05-1997
		AU 2479492 A	16-03-1993
		DE 69219136 D	22-05-1997
		DE 69219136 T	28-08-1997
		DK 601001 T	20-10-1997
		EP 0601001 A	15-06-1994
		ES 2101866 T	16-07-1997
		GR 3023848 T	30-09-1997
		JP 6510286 T	17-11-1994
WO 9609823 A	04-04-1996	AU 3731895 A	19-04-1996
		AU 3731995 A	19-04-1996
		WO 9610015 A	04-04-1996
		US 5628999 A	13-05-1997
		US 5753790 A	19-05-1998
		US 5629319 A	13-05-1997